



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N 37/18, A61K 38/00, C07K 1/00, 2/00, 4/00, 7/00, 14/00, 16/00, 17/00, C07H 21/02, 21/04	A1	(11) International Publication Number: WO 97/48275 (43) International Publication Date: 24 December 1997 (24.12.97)
(21) International Application Number: PCT/US97/10942 (22) International Filing Date: 19 June 1997 (19.06.97) (30) Priority Data: 60/020,150 20 June 1996 (20.06.96) US 08/878,474 18 June 1997 (18.06.97) US (71) Applicant: THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; 22nd floor, 300 Lakeside Drive, Oakland, CA 94612 (US). (72) Inventors: DE ROBERTIS, Edward, M.; 16958 Dulce Ynez Lane, Pacific Palisades, CA 90272 (US). BOUWMEESTER, Tewis; Apartment 708, 827 Levering Avenue, Los Angeles, CA 90024 (US). (74) Agents: SIEBERT, J., Suzanne et al.; Majestic, Parsons, Siebert & Hsue, Suite 1100, Four Embarcadero Center, San Francisco, CA 94111 (US).		(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: ENDODERM, CARDIAC AND NEURAL INDUCING FACTORS (57) Abstract <p>Novel proteins have been designated "cerberus" and "frzb-1", respectively. Cerberus is expressed as a secreted peptide during embryogenesis of the <i>Xenopus</i> embryo, and is expressed specifically in the head organizer region. This new molecule has endodermal, cardiac, and neural tissue inducing activity, that should prove useful in therapeutic, diagnostic, and clinical applications requiring regeneration, differentiation, or repair of these and other tissues. Frzb-1 is a soluble antagonist of growth factors of the Wnt family that acts by binding to Wnt growth factors in the extracellular space. A third novel protein is termed PAPC which promotes the formation of dorsal mesoderm and somites in the embryo.</p>		

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

ENDODERM, CARDIAC AND
NEURAL INDUCING FACTORS

5 Field of the Invention

 The invention generally relates to growth factors, neurotrophic factors, and their inhibitors, and more particularly to several new growth factors with neural, endodermal, and cardiac tissue inducing
10 activity, to complexes and compositions including the factors, and to DNA or RNA coding sequences for the factors. Further, one of the novel growth factors should be useful in tumor suppression gene therapy.

 This application claims the benefit of U.S.
15 Provisional Application No. 60/020,150, filed June 20, 1996.

 This invention was made with Government support under grant contract number HD-21502, awarded by the National Institutes of Health. The Government has
20 certain rights in this invention.

Background of the Invention

 Growth factors are substances, such as polypeptide hormones, which affect the growth of defined populations of animal cells *in vivo* or *in vitro*, but
25 which are not nutrient substances. Proteins involved in the growth and differentiation of tissues may promote or inhibit growth, and promote or inhibit differentiation, and thus the general term "growth factor" includes cytokines, trophic factors, and their inhibitors.

Widespread neuronal cell death accompanies normal development of the central and peripheral nervous systems. Studies of peripheral target tissues during development have shown that neuronal cell death results from the competition among neurons for limiting amounts of survivor factors ("neurotrophic factors"). The earliest identified of these, nerve growth factor ("NGF"), is the most fully characterized and has been shown to be essential for the survival of sympathetic and neural crest-derived sensory neurons during early development of both chick and rat.

One family of neurotropic factors are the Wnts, which have dorsal axis-inducing activity. Most of the Wnt proteins are bound to cell surfaces. (See, e.g., Sokol et al., *Science*, 249, pp. 561-564, 1990.) Dorsal axis-inducing activity in *Xenopus* embryos by one member of this family (Xwnt-8) was described by Smith and Harland in 1991, *Cell*, 67, pp. 753-765. The authors described using RNA injections as a strategy for identifying endogenous RNAs involved in dorsal patterning to rescue dorsal development in embryos that were ventralized by UV irradiation.

Another member of the growth and neurotropic factor family was subsequently discovered and described by Harland and Smith, which they termed "noggin." (*Cell*, 70, pp. 829-840 (1992).) Noggin is a good candidate to function as a signaling molecule in Nieuwkoop's center, by virtue of its maternal transcripts, and in Spemann's organizer, through its zygotic organizer-specific expression. Besides noggin, other secreted factors may be involved in the organizer phenomenon.

Another *Xenopus* gene designated "chordin" that begins to be expressed in Spemann's organizer and that can completely rescue axial development in ventralized

embryos was described by Sasai et al., *Cell*, 79, pp. 779-790, 1994. In addition to dorsalizing mesoderm, chordin has the ability to induce neural tissue and its activities are antagonized by Bone Morphogenetic Protein-4 (Sasai et al., *Nature*, 376, pp. 333-336, 1995).

Therefore, the dorsal lip or Spemann's organizer of the *Xenopus* embryo is an ideal tissue for seeking novel growth and neurotrophic factors. New growth and neurotrophic factors are useful agents, particularly those that are secreted due to their ability to be used in physiologically active, soluble forms because these factors, their receptors, and DNA or RNA coding sequences therefore and fragments thereof are useful in a number of therapeutic, clinical, research, diagnostic, and drug design applications.

Summary of the Invention

In one aspect of the present invention, the sequence of the novel peptide that can be in substantially purified form is shown by SEQ ID NO:1. The *Xenopus* derived SEQ ID NO:1 has been designated "cerberus," and this peptide is capable of inducing endodermal, cardiac, and neural tissue development in vertebrates when expressed. The nucleotide sequence which, when expressed results in cerberus, is illustrated by SEQ ID NO:2. Since peptides of the invention induce endodermal, cardiac, and neural tissue differentiation in vertebrates, they should be able to be prepared in physiologically active form for a number of therapeutic, clinical, and diagnostic applications.

Cerberus was isolated during a search for molecules expressed specifically in Spemann's organizer containing a secretory signal sequence. In addition to cerberus, two other novel cDNAs were identified.

The *Xenopus* derived peptide that can be deduced from SEQ ID NO:3 encodes a novel protein we had earlier designated as "frazzled," a secreted protein of 318 amino acids that has dorsalizing activity in *Xenopus* embryos. We now designate the novel protein as "frzb-1." The gene for frzb-1 is expressed in many adult tissues of many animals, three of the cDNAs (*Xenopus*, mouse, and human) have been cloned by us. The accession numbers for the *Xenopus*, mouse, and human frzb-1 cDNA sequences of the gene now designated frzb-1 are U68059, U68058, and U68057, respectively. Frzb-1 has some degree of sequence similarity to the *Drosophila* gene frizzled which has been shown to encode a seven-transmembrane protein that can act both as a signalling and as a receptor protein (Vinson et al., *Nature*, 338, pp. 263-264, 1989; Vinson and Adler, *Nature*, 329, pp. 549-551, 1987). Vertebrate homologues of Frizzled have been isolated and they too were found to be anchored to the cell membrane by seven membrane spanning domains (Wang et al., *J. Biol. Chem.*, 271, pp. 4468-4476, 1996). Frzb-1 differs from the frizzled proteins in that it is an entirely soluble, diffusible secreted protein and therefore suitable as a therapeutic agent. The nucleotide sequence derived from *Xenopus* that, when expressed, results in frzb-1 protein is illustrated by SEQ ID NO:4. The frzb-1 protein derived from mouse is shown as SEQ ID NO:7, while the mouse frzb-1 nucleotide sequence is SEQ ID NO:8. The human derived frzb-1 protein is illustrated by SEQ ID NO:9, and the human frzb-1 nucleotide sequence is SEQ ID NO:10.

Frzb-1 is an antagonist of Wnts *in vivo*, and thus is believed to find utility as a tumor suppressor gene, since overexpressed Wnt proteins cause cancer. Frzb-1 may also be a useful vehicle for solubilization

and therapeutic delivery of Wnt proteins complexed with it.

The final cDNA isolated containing a signal sequence results in a peptide designated Paraxial
5 Protocadherin (PAPC). The cDNA for PAPC is a divergent member of the cadherin multigene family. PAPC is most related to protocadherin 43 reported by Sano et al., *The EMBO J.*, 12, pp. 2249-2256, 1993. As shown in SEQ ID NO:5, the PAPC gene encodes a transmembrane protein of
10 896 amino acids, of which 187 are part of an intracellular domain. PAPC is a cell adhesion molecule, and microinjection of PAPC mRNA constructs into *Xenopus* embryos suggest that PAPC acts as a molecule involved in mesoderm differentiation. A soluble form of the PAPC
15 extracellular domain is able to block muscle and mesoderm formation in *Xenopus* embryos. The nucleotide sequence encoding *Xenopus* PAPC is provided in SEQ ID NO:6.

Cerberus, frzb-1, or PAPC or fragments thereof
20 (which also may be synthesized by *in vitro* methods) may be fused (by recombinant expression or *in vitro* covalent methods) to an immunogenic polypeptide and this, in turn, may be used to immunize an animal in order to raise antibodies against the novel proteins. Antibodies
25 are recoverable from the serum of immunized animals. Alternatively, monoclonal antibodies may be prepared from cells from the immunized animal in conventional fashion. Immobilized antibodies are useful particularly in the diagnosis (*in vitro* or *in vivo*) or purification
30 of cerberus, frzb-1, or PAPC.

Substitutional, deletional, or insertional mutants of the novel polypeptides may be prepared by *in vitro* or recombinant methods and screened for immuno-crossreactivity with cerberus, frzb-1, or PAPC and for
35 cerberus antagonist or agonist activity.

Cerberus or frzb-1 also may be derivatized in vitro in order to prepare immobilized and labelled proteins, particularly for purposes of diagnosis of insufficiencies thereof, or for affinity purification of antibodies thereto.

Among applications for the novel proteins are tissue replacement therapy and, because frzb-1 is an antagonist of Wnt signaling, tumor suppression therapies. The cerberus receptor may define a novel signalling pathway. In addition, frzb-1 could permit the isolation of novel members of the Wnt family of growth factors.

Brief Description of the Drawings

Figure 1 illustrates the amino acid sequence (SEQ ID NO:1) of the Fig. 2 cDNA clone for cerberus;

Figure 2 illustrates a cDNA clone (SEQ ID NO:2) for cerberus derived from Xenopus. Sense strand is on top (5' to 3' direction) and the antisense strand on the bottom line (in the opposite direction);

Figures 3 and 4 show the amino acid and nucleotide sequence, respectively, of full-length frzb-1 from Xenopus (SEQ ID NOS:3 and 4);

Figures 5 and 6 show the amino acid and nucleotide sequence, respectively, of full-length PAPC from Xenopus (SEQ ID NOS:5 and 6);

Figures 7 and 8 show the amino acid and nucleotide sequence, respectively, of full-length frzb-1 from mouse (SEQ ID NOS:7 and 8); and

Figures 9 and 10 show the amino acid and nucleotide sequence, respectively, of full-length frzb-1 from human (SEQ ID NOS:9 and 10).

Detailed Description of the Preferred Embodiments

Among the several novel proteins and their nucleotide sequences described herein, is a novel endodermal, cardiac, and neural inducing factor in vertebrates that we have named "cerberus." When referring to cerberus, the present invention also contemplates the use of fragments, derivatives, agonists, or antagonists of cerberus molecules. Because cerberus has no homology to any reported growth factors, it is proposed to be the founding member of a novel family of growth factors with potent biological activities, which may be isolated using SEQ ID NO:2.

The amphibian organizer consists of several cell populations with region-specific inducing activities. On the basis of morphogenetic movements, three very different cell populations can be distinguished in the organizer. First, cells with crawling migration movements involute, fanning out to form the prechordal plate. Second, cells involute through the dorsal lip driven by convergence and extension movements, giving rise to the notochord of the trunk. Third, involution ceases and the continuation of mediolateral intercalation movements leads to posterior extension movements and to the formation of the tail notochord and of the chordoneural hinge. The three cell populations correspond to the head, trunk, and tail organizers, respectively.

The cerberus gene is expressed at the right time and place to participate in cell signalling by Spemann's organizer. Specifically, cerberus is expressed in the head organizing region that consists of crawling-migrating cells. The cerberus expressing region corresponds to the prospective foregut, including the liver and pancreas anlage, and the heart mesoderm.

Cerberus expression is activated by chordin, noggin, and organizer-specific homeobox genes.

Our studies were conducted in early embryos of the frog *Xenopus laevis*. The frog embryo is well suited to experiments, particularly experiments pertaining to generating and maintaining regional differences within the embryo for determining roles in tissue differentiation. It is easy to culture embryos with access to the embryos even at very early stages of development (preceding and during the formation of body pattern and differentiation) and the embryos are large. The initial work with noggin and chordin also had been in *Xenopus* embryos, and, as predicted, was highly conserved among vertebrates. Predictions based on work with *Xenopus* as to corresponding human noggin were proven true and the ability to clone the gene for human noggin was readily accomplished. (See the description of *Xenopus* work and cloning information in PCT application, published March 17, 1994, WO 9 405 800, and the subsequent human cloning based thereon in the PCT application, also published March 17, 1994, as WO 9 405 791.)

CLONING

The cloning of cerberus, frzb-1, and PAPC resulted from a comprehensive screen for cDNAs enriched in Spemann's organizer. Subtractive differential screening was performed as follows. In brief, poly A⁺ RNA was isolated from 300 dorsal lip and ventral marginal zone (VMZ) explants at stage 10½. After first strand cDNA synthesis approximately 70-80% of common sequences were removed by subtraction with biotinylated VMZ poly A⁺ RNA prepared from 1500 ventral gastrula halves. For differential screening, duplicate filters (2000 plaques per 15 cm plate, a total of 80,000 clones

screened) of an unamplified oriented dorsal lip library were hybridized with radiolabeled dorsal lip or VMZ cDNA. Putative organizer-specific clones were isolated, grouped by sequence analysis from the 5' end and whole-mount in situ hybridization, and subsequently classified into known and new dorsal-specific genes. Rescreening of the library (100,000 independent phages) with a cerberus probe resulted in the isolation of 45 additional clones, 31 of which had similar size as the longest one of the 11 original clones indicating that they were presumably full-length cDNAs. The longest cDNAs for cerberus, frzb-1, and PAPC were completely sequenced.

To explore the molecular complexity of Spemann's organizer we performed a comprehensive differential screen for dorsal-specific cDNAs. The method was designed to identify abundant cDNAs without bias as to their function. As shown in Table 1, five previously known cDNAs and five new ones were isolated, of which three (expressed as cerberus, frzb-1, and PAPC, respectively) had secretory signal sequences.

TABLE 1

	Previously Known Genes	Gene Product	No. of Isolates
	Chordin	novel secreted protein	70
	Goosecoid	homeobox gene	3
5	Pintallavis/XFKH-1	forkhead/transcription factor	2
	Xnot-2	homeobox gene	1
	Xlim-1	homeobox gene	1
	New Genes		
	Cerberus	novel secreted protein	11
10	PAPC	cadherin-like/transmembrane	2
	Frzb-1	novel secreted protein	1
	Sox-2	sry/transcription factor	1
	Fkh-like	forkhead/transcription factor	1

15 The most abundant dorsal-specific cDNA was
 chordin (chd), with 70 independent isolates. The second
 most abundant cDNA was isolated 11 times and named
 cerberus (after a mythological guardian dog with
 multiple heads). The cerberus cDNA encodes a putative
 20 secreted polypeptide of 270 amino acids, with an amino
 terminal hydrophobic signal sequence and a carboxy
 terminal cysteine-rich region (Fig. 1). Cerberus is
 expressed specifically in the head organizer region of
 the *Xenopus* embryo, including the future foregut.

25 An abundant mRNA found in the dorsal region of
 the *Xenopus* gastrula encodes the novel putative secreted
 protein we have designated as cerberus. Cerberus mRNA
 has potent inducing activity in *Xenopus* embryos, leading
 to the formation of ectopic heads. Unlike other
 organizer-specific factors, cerberus does not dorsalize
 30 mesoderm and is instead an inhibitor of trunk-tail
 mesoderm. Cerberus is expressed in the anterior-most

domain of the gastrula including the leading edge of the deep layer of the dorsal lip a region that, as shown here, gives rise to foregut and midgut endoderm. Cerberus promotes the formation of cement gland, olfactory placodes, cyclopic eyes, forebrain, and duplicated heart and liver (a foregut derivative). Because the pancreas is also derived from this foregut region, it is likely that cerberus induces pancreas in addition to liver. The expression pattern and inducing activities of cerberus suggest a role for a previously neglected region of the embryo, the prospective foregut endoderm, in the induction of the anterior head region of the embryo.

Turning to Fig. 1, *Xenopus cerberus* encodes a putative secreted protein transiently expressed during embryogenesis and the deduced amino acid sequence of *Xenopus cerberus* is shown. The signal peptide sequence and the nine cysteine residues in the carboxy-terminus are indicated in bold. Potential N-linked glycosylation sites are underlined. In database searches the cerberus protein showed limited similarity only to the mammalian Dan protein, a possible tumor suppressor proposed to be a DNA-binding protein.

Cerberus appears to be a pioneer protein, as its amino acid sequence and the spacing of its 9 cysteine residues were not significantly similar to other proteins in the databases (NCBI-Gen Bank release 93.0). We conclude that the second most abundant dorsal-specific cDNA encodes a novel putative secreted factor, which should be the founding member of a novel family of growth factors active in cell differentiation.

Cerberus Demarcates an Anterior Organizer Domain. Cerberus mRNA is expressed at low levels in the unfertilized egg, and zygotic transcripts start accumulating at early gastrula. Expression continues

during gastrula and early neurula, rapidly declining during neurulation. Importantly, cerberus expression starts about one hour after that of chd, suggesting that cerberus could act downstream of the chd signal.

5 Whole-mount *in situ* hybridizations reveal that expression starts in the yolky endomesodermal cells located in the deep layer of the organizer. The cerberus domain includes the leading edge of the most anterior organizer cells and extends into the lateral
10 mesoderm. The leading edge gives rise to liver, pancreas, and foregut in its midline, and the more lateral region gives rise to heart mesoderm at later stages of development.

Fig. 2 sets out the sequence of a full length
15 *Xenopus* cDNA for cerberus.

This entirely new molecule has demonstrated physiological properties that should prove useful in therapeutic, diagnostic, and clinical applications that require regeneration, differentiation, or repair of
20 tissues, such wound repair, neuronal regenerative or transplantation, supplementation of heart muscle differentiation, differentiation of pancreas and liver, and other applications in which cell differentiation processes are to be induced.

25 The second, novel, secreted protein we have discovered is called "frzb-1," which was shown to be a secreted protein in *Xenopus* oocyte microinjection experiments. Thus it provides a natural soluble form of the related extracellular domains of *Drosophila* and
30 vertebrate frizzled proteins. We propose that the latter proteins could be converted into active soluble forms by introducing a stop codon before the first transmembrane domain. We have noted that the cysteine-rich region of frzb-1 and frizzled contains some overall
35 structural homology with Wnt proteins using the Profile

Search homology program (Gribskov, *Meth. Enzymol.*, 183, pp. 146-159, 1990). This had raised the interesting possibility that frzb-1 could interact directly with Wnt growth factors in the extracellular space. This was

5 because we had found that when microinjected into *Xenopus* embryos, frzb-1 constructs have moderate dorsalizing activity, leading to the formation of embryos with enlarged brain and head, and shortened trunk. Somatic muscle differentiation, which requires

10 Xwnt-8, was inhibited. In the case of frzb-1, an attractive hypothesis, suggested by the structural homologies, was that it may act as an inhibitor of Wnt-8, a growth factor that has ventralizing activity in the *Xenopus* embryo (Christian and Moon, *Genes Dev.*, 7,

15 pp. 13-28, 1993). We have shown that frzb-1 can interact with Xwnt-8 and Wnt-1, and it is expected that it could also interact with other members of the Wnt family of growth factors, of which at least 15 members exist in mammals. In addition, a possible interaction

20 with Wnts was suggested by the recent discovery that dishevelled, a gene acting downstream of wingless, has strong genetic interaction with frizzled mutants in *Drosophila* (Krasnow et al., *Development*, 121, pp. 4095-4102, 1995). This possibility has been explored in

25 depth (Leyns et al., *Cell*, 88, pp. 747-756, March 21, 1997), because a soluble antagonist of the Wnt family of proteins is expected to be of great therapeutic value. Examples 1 and 2 illustrate tests that show antagonism of Xwnt-8 by binding to frzb-1.

30 Vertebrate homologues of Frizzled have been isolated and they too are anchored to the cell membrane by seven membrane spanning domains (Wang et al., *J. Biol. Chem.*, 271, pp. 4468-4476, 1996). Frzb-1 differs from the frizzled proteins in that it is an

35 entirely soluble, diffusible secreted protein and

therefore suitable as a therapeutic agent. The nucleotide sequence that when expressed results in frzb-1 protein is illustrated by SEQ ID NO:4.

SEQ ID NO:4 corresponds to the *Xenopus* homolog, but by using it in BLAST searches (and by cloning mouse frzb-1) we had been able to assemble the sequence of the entire mature human frzb-1 protein, SEQ ID NO:9. Indeed, human frzb-1 is encoded in six expressed sequence tags (ESTs) available in Genbank. The human frzb-1 sequence can be assembled by overlapping in the 5' to 3' direction the ESTs with the following accession numbers in Genbank: H18848, R63748, W38677, W44760, H38379, and N71244. No function had yet been assigned to these EST sequences, but we believe and thus propose here that human frzb-1 will have similar functions in cell differentiation to those described above for *Xenopus* frzb-1. The nucleotide sequence of human frzb-1 is shown in SEQ ID NO:10. The mouse frzb-1 protein and nucleotide sequences are provided by SEQ ID NOS:7 and 8, respectively.

In particular, we believe that frzb-1 will prove useful in gene therapy of human cancer cells. In this rapidly developing field, one approach is to introduce vectors expressing anti-sense sequences to block expression of dominant oncogenes and growth factor receptors. Another approach is to produce episomal vectors that will replicate in human cells in a controlled fashion without transforming the cells. For an example of the latter (an episomal expression vector system for human gene therapy), reference is made to U.S. Patent 5,624,820, issued April 29, 1997, inventor Cooper.

Gene therapy now includes uses of human tumor suppression genes. For example, U.S. Patent 5,491,064, issued February 13, 1996, discloses a tumor suppression

gene localized on chromosome 11 and described as potentially useful for gene therapy in cancers deleted or altered in their expression of that gene. Frzb-1 maps to chromosome 2q31-33 and loss of one copy of the
5 2q31-33 and loss of one copy of the 2q arm has been observed with high incidence in lung carcinomas, colo-rectal carcinomas, and neuroblastomas, which has lead to the proposal that the 2q arm carries a tumor suppressor gene. We expect frzb to be a tumor
10 suppressor gene, and thus to be useful in tumor suppression applications.

A number of applications for cerberus and frzb-1 are suggested from their pharmacological (biological activity) properties.

15 For example, the cerberus and frzb-1 cDNAs should be useful as a diagnostic tool (such as through use of antibodies in assays for proteins in cell lines or use of oligonucleotides as primers in a PCR test to amplify those with sequence similarities to the
20 oligonucleotide primer, and to determine how much of the novel protein is present).

Cerberus, of course, might act upon its target cells via its own receptor. Cerberus, therefore, provides the key to isolate this receptor. Since many
25 receptors mutate to cellular oncogenes, the cerberus receptor should prove useful as a diagnostic probe for certain tumor types. Thus, when one views cerberus as ligand in complexes, then complexes in accordance with the invention include antibody bound to cerberus,
30 antibody bound to peptides derived from cerberus, cerberus bound to its receptor, or peptides derived from cerberus bound to its receptor or other factors. Mutant forms of cerberus, which are either more potent agonists or antagonists, are believed to be clinically useful.

Such complexes of cerberus and its binding protein partners will find uses in a number of applications.

Practice of this invention includes use of an oligonucleotide construct comprising a sequence coding
5 for cerberus or frzb-1 and for a promoter sequence
operatively linked in a mammalian or a viral expression
vector. Expression and cloning vectors contain a
nucleotide sequence that enables the vector to replicate
in one or more selected host cells. Generally, in
10 cloning vectors this sequence is one that enables the
vector to replicate independently of the host
chromosomes, and includes origins of replication or
autonomously replicating sequences. The well-known
plasmid pBR322 is suitable for most gram negative
15 bacteria, the 2 μ plasmid origin for yeast and various
viral origins (SV40, polyoma, adenovirus, VSV or BPV)
are useful for cloning vectors in mammalian cells.

Expression and cloning vectors should contain
a selection gene, also termed a selectable marker.
20 Typically, this is a gene that encodes a protein
necessary for the survival or growth of a host cell
transformed with the vector. The presence of this gene
ensures that any host cell which deletes the vector will
not obtain an advantage in growth or reproduction over
25 transformed hosts. Typical selection genes encode
proteins that (a) confer resistance to antibiotics or
other toxins, e.g. ampicillin, neomycin, methotrexate or
tetracycline, (b) complement auxotrophic deficiencies.

Examples of suitable selectable markers for
30 mammalian cells are dihydrofolate reductase (DHFR) or
thymidine kinase. Such markers enable the identifica-
tion of cells which were competent to take up the
cerberus nucleic acid. The mammalian cell transformants
are placed under selection pressure which only the
35 transformants are uniquely adapted to survive by virtue

of having taken up the marker. Selection pressure is imposed by culturing the transformants under conditions in which the concentration of selection agent in the medium is successively changed. Amplification is the process by which genes in greater demand for the production of a protein critical for growth are reiterated in tandem within the chromosomes of successive generations of recombinant cells. Increased quantities of cerberus or frzb-1 can therefor be synthesized from the amplified DNA.

For example, cells transformed with the DHFR selection gene are first identified by culturing all of the transformants in a culture medium which contains methotrexate (Mtx), a competitive antagonist of DHFR. An appropriate host cell in this case is the Chinese hamster ovary (CHO) cell line deficient in DHFR activity, prepared and propagated as described by Urlaub and Chasin, *Proc. Nat. Acad. Sci.*, 77, 4216 (1980). The transformed cells then are exposed to increased levels of Mtx. This leads to the synthesis of multiple copies of the DHFR gene and, concomitantly, multiple copies of other DNA comprising the expression vectors, such as the DNA encoding cerberus or frzb-1. Alternatively, host cells transformed by an expression vector comprising DNA sequences encoding cerberus or frzb-1 and aminoglycoside 3' phosphotransferase (APH) protein can be selected by cell growth in medium containing an aminoglycosidic antibiotic such as kanamycin or neomycin or G418. Because eukaryotic cells do not normally express an endogenous APH activity, genes encoding APH protein, commonly referred to as neo resistant genes, may be used as dominant selectable markers in a wide range of eukaryotic host cells, by which cells transformed by the vector can readily be identified.

Expression vectors, unlike cloning vectors, should contain a promoter which is recognized by the host organism and is operably linked to the cerberus nucleic acid. Promoters are untranslated sequences located upstream from the start codon of a structural gene (generally within about 100 to 1000 bp) that control the transcription and translation of nucleic acid under their control. They typically fall into two classes, inducible and constitutive. Inducible promoters are promoters that initiate increased levels of transcription from DNA under their control in response to some change in culture conditions, e.g. the presence or absence of a nutrient or a change in temperature. At this time a large number of promoters recognized by a variety of potential host cells are well known. These promoters can be operably linked to cerberus encoding DNA by removing them from their gene of origin by restriction enzyme digestion, followed by insertion 5' to the start codon for cerberus or frzb-1.

Nucleic acid is operably linked when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein which participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, operably linked means that the DNA sequences being linked are contiguous and, in the case of a secretory leader, contiguous and in reading phase. Linking is accomplished by ligation at convenient restriction sites. If such sites do not

exit then synthetic oligonucleotide adapters or linkers are used in accord with conventional practice.

Transcription of the protein-encoding DNA in mammalian host cells is controlled by promoters obtained from the genomes of viruses such as polyoma, cytomegalovirus, adenovirus, retroviruses, hepatitis-B virus, and most preferably Simian Virus 40 (SV40), or from heterologous mammalian promoters, e.g. the actin promoter. Of course, promoters from the host cell or related species also are useful herein.

Cerberus and frzb-1 are clearly useful as a component of culture media for use in culturing cells, such as endodermal, cardiac, and nerve cells, *in vitro*. We believe cerberus and frzb-1 will find uses as agents for enhancing the survival or inducing the growth of liver, pancreas, heart, and nerve cells, such as in tissue replacement therapy.

The final cDNA isolated containing a signal sequence results in a peptide designated Paraxial Protocadherin (PAPC). The cDNA for PAPC is a divergent member of the cadherin multigene family. PAPC is most related to protocadherin 43 reported by Sano et al., *The EMBO J.*, 12, pp. 2249-2256, 1993. As shown in SEQ ID NO:5, the PAPC gene encodes a transmembrane protein of 896 amino acids, of which 187 are part of an intracellular domain. PAPC is a cell adhesion molecule, and microinjection of PAPC mRNA constructs into *Xenopus* embryos suggest that PAPC acts in mesoderm differentiation. The nucleotide sequence encoding *Xenopus* PAPC is provided in SEQ ID NO:6.

Therapeutic formulations of the novel proteins may be prepared for storage by mixing the polypeptides having the desired degree of purity with optional physiologically acceptable carriers, excipients or stabilizers, in the form of lyophilized cake or aqueous

solutions. Acceptable carriers, excipients or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, and other organic acids; anti-oxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin or immunoglobulins. Other components can include glycine, glutamine, asparagine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as Tween, Pluronic or PEG.

Polyclonal antibodies to the novel proteins generally are raised in animals by multiple subcutaneous (sc) or intraperitoneal (ip) injections of cerberus or frzb-1 and an adjuvant. It may be useful to conjugate these proteins or a fragment containing the target amino acid sequence to a protein which is immunogenic in the species to be immunized, e.g., keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, or soybean trypsin inhibitor using a bifunctional or derivatizing agent, for example, maleimidobenzoyl sulfosuccinimide ester (conjugation through cysteine residues), N-hydroxysuccinimide (through lysine residues), glutaraldehyde, succinic anhydride, SOCl_2 , or $\text{R}^1\text{N} = \text{C} = \text{NR}$.

Animals can be immunized against the immunogenic conjugates or derivatives by combining 1 mg or 1 μg of conjugate (for rabbits or mice, respectively) with 3 volumes of Freund's complete adjuvant and injecting the solution intradermally in multiple sites. One month later the animals are boosted with 1/5 to 1/10 the original amount of conjugate in Freund's complete

adjuvant by subcutaneous injection at multiple sites. Seven to 14 days later animals are bled and the serum is assayed for anti-cerberus titer. Animals are boosted until the titer plateaus. Preferably, the animal is
5 boosted with the conjugate of the same cerberus or frzb-1 polypeptide, but conjugated to a different protein and/or through a different cross-linking agent. Conjugates also can be made in recombinant cell culture as protein fusions. Also, aggregating agents such as
10 alum are used to enhance the immune response.

Monoclonal antibodies are prepared by recovering spleen cells from immunized animals and immortalizing the cells in conventional fashion, e.g. by fusion with myeloma cells or by EB virus transformation
15 and screening for clones expressing the desired antibody.

Antibodies are useful in diagnostic assays for cerberus, frzb-1, or PAPC or their antibodies and to identify family members. In one embodiment of a
20 receptor binding assay, an antibody composition which binds to all of a selected plurality of members of the cerberus family is immobilized on an insoluble matrix, the test sample is contacted with the immobilized antibody composition in order to adsorb all cerberus
25 family members, and then the immobilized family members are contacted with a plurality of antibodies specific for each member, each of the antibodies being individually identifiable as specific for a predetermined family member, as by unique labels such as
30 discrete fluorophores or the like. By determining the presence and/or amount of each unique label, the relative proportion and amount of each family member can be determined.

The antibodies also are useful for the
35 affinity purification of the novel proteins from

recombinant cell culture or natural sources. Antibodies that do not detectably cross-react with other growth factors can be used to purify the proteins free from these other family members.

5

EXAMPLE 1

Frzb-1 Antagonizes Xwnt-8 Non-Cell Autonomously

To test whether frzb-1 can antagonize secondary axes caused by Xwnt-8 after secretion by injected cells, an experimental design was used. Thus, frzb-1 mRNA was injected into each of the four animal blastomeres of eight-cell embryos, and subsequently, a single injection of Xwnt-8 mRNA was given to a vegetal-ventral blastomere at the 16-32 cell stage. In two independent experiments, we found that injection of frzb-1 alone (n=13) caused mild dorsalization with enlargement of the cement gland in all embryos and that injection of Xwnt-8 alone (n=53) lead to induction of complete secondary axes in 67% of the embryos. However, injection of frzb-1 into animal caps abolished the formation of complete axes induced by Xwnt-8 (n=27), leaving only a residual 14% of embryos with very weak secondary axes. The double-injected embryos retained the enlarged cement gland phenotype caused by injection of frzb-1 mRNA alone. Because both mRNAs encode secreted proteins and were microinjected into different cells, we conclude that the antagonistic effects of frzb-1 and Xwnt-8 took place in the extracellular space after these proteins were secreted.

EXAMPLE 2

Membrane-Anchored Wnt-1 Confers Frzb-1 Binding

To investigate a possible interaction between frzb-1 and Wnts, the first step was to insert an HA epitope tag into a Xenopus frzb-1 construct driven by the CMV (cytomegalovirus) promoter. Frzbl-HA was tested in mRNA microinjection assays in Xenopus embryos and found to be biologically active. Conditioned medium from transiently transfected cells contained up to 10 $\mu\text{g/ml}$ of Frzbl-HA (quantitated on Western blots using an HA-tagged protein standard).

Transient transfection of 293 cells has been instrumental in demonstrating interactions between wingless and frizzled proteins. We therefore took advantage of constructs in which Wnt-1 was fused at the amino terminus of CD8, generating a transmembrane protein containing biologically active Wnt-1 exposed to the extracellular compartment. A Wnt1CD8 cDNA construct (a generous gift of Dr. H. Varmus, NIH) was subcloned into the pCDNA (Invitrogen) vector and transfected into 293 cells. After incubation with Frzbl-HA-conditioned medium (overnight at 37°C), intensely labeled cells were observed by immunofluorescence. As a negative control, a construct containing 120 amino acids of Xenopus chordin, an unrelated secreted protein was used. Transfection of this construct produced background binding of Frzbl-HA to the extracellular matrix, both uniform and punctate. Cotransfection of Wnt1CD8 with pCDNA-LacZ showed that transfected cells stained positively for Frzbl-HA and LacZ. Since Wnt1CD8 contains the entire CD8 molecule, a CD8 cDNA was used as an additional negative control. After transfection with LacZ and full-length CE8, Frzbl-HA failed to bind to the transfected cells. Although most of our experiments

were carried out at 37°C, Frzb1-HA-conditioned medium also stained Wnt1CD8-transfected cells after incubation at 4°C for 2 hours.

5 Attempts to biochemically quantitate the binding of Frzb-1 to Wnt1CD8-transfected cells were unsuccessful due to high background binding to control cultures, presumably due to binding to the extracellular matrix. Thus, we were unable to estimate a K_D for the affinity of the Frzb-1/Wnt-1 interaction. However, when
10 serial dilutions of conditioned medium containing Frzb1-HA were performed (ranging from 2.5×10^{-7} to 1.25×10^{-10} M), staining of Wnt1CD8-transfected cells was found at all concentrations.

15 Although we have been unable to provide biochemical evidence for direct binding between Wnts and frzb-1, this cell biological assay indicates that Frzb1-HA can bind, directly or indirectly, to Wnt-1 on the cell membrane in the 10^{-10} M range.

20 It is to be understood that while the invention has been described above in conjunction with preferred specific embodiments, the description and examples are intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims.

It is Claimed:

1. A substantially pure protein characterized by a physiologically active form and comprising an amino acid sequence encoded by the DNA of SEQ ID NO:2.
2. The protein as in claim 1 having neurotrophic, growth or differentiation factor activity.
3. A composition comprising the protein of claim 1 and a physiologically acceptable carrier with which the peptide is admixed.
4. An oligonucleotide construct comprising a sequence coding for a protein and an expression vector operatively linked therewith, the protein having neurotrophic, growth or differentiation factor activity
5 and being expressible from SEQ ID NO:2.
5. The construct as in claim 4 wherein the expression vector is a mammalian or viral expression vector.
6. A substantially pure protein characterized by a physiologically active form and comprising an amino acid sequence encoded by the DNA of SEQ ID NO:4, SEQ ID NO:8, or SEQ ID NO:10.
7. The protein as in claim 6 having neurotrophic, growth or differentiation factor activity.
8. A composition comprising the protein of claim 6 and a physiologically acceptable carrier with which the protein is admixed.

9. An oligonucleotide construct comprising a sequence coding for a protein and an expression vector operatively linked therewith, the protein being expressible from SEQ ID NO:4, SEQ ID NO:8 or SEQ ID NO:10.

10. The construct as in claim 9 wherein the protein is expressible in soluble form.

11. The construct as in claim 9 wherein the expression vector is a mammalian or viral expression vector.

12. A complex comprising a substantially pure frzb-1 protein complexed with at least one Wnt protein.

13. A substantially pure protein characterized by a physiologically active form and comprising an amino acid sequence encoded by the DNA of SEQ ID NO:6.

14. The protein as in claim 13 having mesoderm differentiation activity.

15. A composition comprising the protein of claim 13 and a physiologically acceptable carrier with which the protein is admixed.

MLLNVLRICI	IVCLVNDGAG	KHSEGRERTK	TYSLNSRGYF	40
RKERGARRSK	ILLVNTKGLD	EPHIGHGDFG	LVAELFDSTR	80
THTNRKEPDM	NKVKLFSTVA	HGNKSARRKA	YNGSRRNIFS	120
RRSFDKRNTE	VTEKPGAKMF	WNNFLVKMNG	APQNTSHGSK	160
AQEIMKEACK	TLPFTQNIVH	ENCDRMVIQN	NLCFGKICISL	200
HVPNQDRRN	TCSHCLPSKF	TLNHLTLNCT	GSKNVVKVVM	240
MVEECTCEAH	KSNFHQTAQF	NMDTSTTLHH		270

Figure 1

SUBSTITUTE SHEET (RULE 26)

GAATTC	CCAG	CAAGTCGCTC	AGAAACACTG	CAGGGTCTAG	ATATCATACA	ATGTTACTAA	60
CTTAAG	GGGTC	GTTCAGCGAG	TCTTTGTGAC	GTCCCAGATC	TATAGTATGT	TACAATGATT	
ATGTACTCAG	GATCTGTATT	ATCGTCTGCC	TTGTGAATGA	TGGAGCAGGA	AAACACTCAG	120	
TACATGAGTC	CTAGACATAA	TAGCAGACGG	AACACTTACT	ACCTCGTCTT	TTTGTGAGTC		
AAGGACGAGA	AAGGACAAAA	ACATATTAC	TTAACAGCAG	AGGTTACTTC	AGAAAAGAAA	180	
TTCTGCTCT	TTCTGTTTT	TGTATAAGTG	AATTGTCGTC	TCCAATGAAG	TCTTTTCTTT		
GAGGAGCAGC	TAGGAGCAAG	ATTCTGCTGG	TGAATACTAA	AGGTCTTGAT	GAACCCACAC	240	
CTCCTCGTGC	ATCCTCGTTC	TAAGACGACC	ACTTATGATT	TCCAGAACTA	CTTGGGGTGT		
TTGGGCATGG	TGATTTTCGC	TTAGTAGCTG	AACTATTGGA	TTCCACCAGA	ACACATACAA	300	
AACCCGTACC	ACTAAAAGCG	AATCATCGAC	TTGATAAACT	AAGGTGGTCT	TGTGTATGTT		
ACAGAAAAGA	GCCAGACATG	AACAAAGTCA	AGCTTTTCTC	AACAGTTGCC	CATGGAAACA	360	
TGTCTTTTCT	CGGTCTGTAC	TTGTTTCAGT	TCGAAAAGAG	TTGTCAACGG	GTACCTTTGT		
AAAGTGCAAG	AAGAAAAGCT	TACAATGGTT	CTAGAAGGAA	TATTTTCTCT	CGCGTTCTT	420	
TTTCAOGTTC	TTCTTTTCTG	ATGTTACCAA	GATCTTCTCT	ATAAAAAGGA	GCGGCAAGAA		
TTGATAAAAG	AAATACAGAG	GTTACTGAAA	AGCCTGGTGC	CAAGATGTTT	TGGAACAATT	480	
AACTATTTTC	TTTATGTCTC	CAATGACTTT	TCGGACCACG	GTTCTACAAG	ACCTTGTTAA		
TTTTGGTTAA	AATGAATGGA	GCCCCACAGA	ATACAAGCCA	TGGCAGTAAA	GCACAGGAAA	540	
AAAACCAATT	TTACTTACCT	CGGGGTGTCT	TATGTTGGGT	ACCGTCATTT	CGTGTCTTTT		
TAATGAAAGA	AGCTTGCAAA	ACCTTGTTTT	TCACTCAGAA	TATTGTACAT	GAAAACGTGT	600	
ATTACTTTCT	TCGAACGTTT	TGGAACAAA	AGTGAGTCTT	ATAACATGTA	CTTTTGACAC		
ACAGGATGGT	GATACAGAAC	AATCTGTGCT	TTGGTAAATG	CATCTCTCTC	CATGTTCCAA	660	
TGTOCTAOC	CTATGTCTTG	TTAGACACGA	AACCATTTAC	GTAGAGAGAG	GTACAAGGTT		
ATCAGCAAGA	TCGACGAAAT	ACTTGTTOCC	ATTGCTTGCC	GTCCAAATTT	ACCCTGAACC	720	
TAGTOGTTCT	AGCTGCTTTA	TGAACAAGGG	TAAOGAACGG	CAGGTTTAAA	TGGGACTTGG		
ACCTGACGCT	GAATTGTACT	GGATCTAAGA	ATGTAGTAAA	GGTTGTCATG	ATGGTAGAGG	780	
TGGACTGOGA	CTTAACATGA	CCTAGATTCT	TACATCATTT	CCAACAGTAC	TACCATCTCC		
AATGCAOGTG	TGAAGCTCAT	AAGAGCAACT	TCCAOCBAAC	TGCACAGTTT	AACATGGATA	840	
TTACGTGCAC	ACTTCGAGTA	TTCTOGTTGA	AGGTGGTTTG	ACGTGTCAAA	TTGTACCTAT		
CATCTACTAC	CCTGCACCAT	TAAAGGACTG	CCATACAGTA	TGGAAATGCC	CTTTTGTGTTG	900	
GTAGATGATG	GGACGTGGTA	ATTCTCTGAC	GGTATGTCAT	ACCTTTACGG	GAAAACAACC		
AAATTTTGT	ACATACTATG	CATCTAAAGC	ATTATGTTGC	CTTCTATTTT	ATATAACCAC	960	
TTATAAACAA	TGTATGATAC	GTAGATTTTC	TAATACAACG	GAAGATAAAG	TATATTGGTG		
ATGGAATAAG	GATTGTATGA	ATTATAATTA	ACAAATGGCA	TTTTGTGTAA	CATGCAAGAT	1020	
TACCTTATTC	CTAACATACT	TAATATTAAT	TGTTTACCGT	AAAACACATT	GTACGTCTTA		

Figure 2A

SUBSTITUTE SHEET (RULE 28)

CTCTGTTCCA	TCAGTTGCAA	GATAAAAGGC	AATATTTGTT	TGACTTTTTT	TCTACAAAAT	1080
GAGACAAGGT	AGTCAACGTT	CTATTTTCCG	TTATAAACAA	ACTGAAAAAA	AGATGTTTAA	
GAATACCCAA	ATATATGATA	AGATAATGGG	GTCAAAACTG	TTAAGGGGTA	ATGTAATAAT	1140
CTTATGGGTT	TATATACTAT	TCTATTACCC	CAGTTTTGAC	AATTCCCAT	TACATTATTA	
AGGGACTAAG	TTTGCCCAGG	AGCAGTGACC	CATAACAACC	AATCAGCAGG	TATGATTTAC	1200
TCCCTGATTG	AAACGGGTCC	TCGTCACTGG	GTATTGTTGG	TTAGTCGTCC	ATACTAAATG	
TGGTCACCTG	TTTAAAAGCA	AACATCTTAT	TGGTTGCTAT	GGGTACTGTC	TTCTGGGCAA	1260
ACCAAGTGGAC	AAATTTTCGT	TTGTAGAATA	ACCAACGATA	CCCAATGACG	AAGACCCGTT	
AATGTGTGCC	TCATAGGGGG	GTTAGTGTGT	TGTGTACTGA	ATAAATTGTA	TTTATTTTAT	1320
TTACACACGG	AGTATCCCCC	CAATCACACA	ACACATGACT	TATTTAACAT	AAATAAAGTA	
TGTTACAAA	AAAAAAA					
ACAATGTTTT	TTTTTTTT					

Figure 2B

MSRTRKVDL LLLAIPGLAL LLLPNAYCAS CEPVRIPMCK SMPWNMTKMP NHLHHSTQAN	60
AILAIEQFEG LLTTECSQDL LFFLCAMYAP ICTIDFQHEP IKPCKSV CER ARAGCEPILI	120
KYRHTWPESL ACEELPVYDR GVCISPEAIV TVEQGTDSMP DFSMDSNNGN CGSGREHCKC	180
KPMKATQKTY LKNNYNYVIR AKVKEVKV KC HDATAIVEVK EILKSSLVNI PKDTVTLYTN	240
SGCLCPQLVA NEEYIIMGYE DKERTRLLLV EGSLAEKWRD RLAKKV KRD QKLRRPRKSK	300
DPVAPIPNKN SNSRQARS	

Figure 3

SUBSTITUTE SHEET (RULE 26)

GAATTCOCTT TCACACAGGA CTCCTGGCAG AGGTGAATGG TTAGCCCTAT GGATTTGGTT	60
CTTAAGGGAA AGTGTGTCCT GAGGACCGTC TCCACTTACC AATCGGGATA CCTAAACCAA	
TGTTGATTTT GACACATGAT TGATTGCTTT CAGATAGGAT TGAAGGACTT GGATTTTAT	120
ACAACTAAAA CTGTGTACTA ACTAACGAAA GTCTATCCTA ACTTCCTGAA CCTAAAAATA	
CTAATTCTGC ACTTTTAAAT TATCTGAGTA ATTGTTTATT TTGTATTGGA TGGGACTAAA	180
GATTAAGACG TGAAAATTTA ATAGACTCAT TAACAAGTAA AACATAACCT ACCCTGATTT	
GATAAACTTA ACTOCTTGCT TTTGACTTGC CCATAAACTA TAAGGTGGGG TGAGTTGTAG	240
CTATTTGAAT TGAGGAACGA AAAGTGAACG GGTATTTGAT ATTCCACCCC ACTCAACATC	
TTGCTTTTAC ATGTGCCAG ATTTTCCCTG TATTCCCTGT ATTCCCTCTA AAGTAAGCCT	300
AACGAAAATG TACACGGGTC TAAAAGGGAC ATAAGGGACA TAAGGGAGAT TTCATTCGGA	
ACACATACAG GTTGGGCAGA ATAACAATGT CTCGAACAAG GAAAGTGGAC TCATTACTGC	360
TGTGTATGTC CAACCCGTCT TATTGTTACA GAGCTTGTTT CTTTCACCTG AGTAATGACG	
TACTGGCCAT ACCTGGACTG GCGCTTCTCT TATTACCCAA TGCTTACTGT GCTTCGTGTG	420
ATGACCGGTA TGGACCTGAC CGCGAAGAGA ATAATGGGTT ACGAATGACA CGAAGCACAC	
AGCCTGTGCG GATCCCCATG TGCAAATCTA TGCCATGGAA CATGACCAAG ATGCCCAACC	480
TCGGACACGC CTAGGGGTAC ACGTTTAGAT ACGGTACCTT GTACTGGTTC TACGGGTTGG	
ATCTCCACCA CAGCACTCAA GCCAATGCCA TCCTGGCAAT TGAACAGTTT GAAGGTTTGC	540
TAGAGGTGGT GTCGTGAGTT CGGTACGGT AGGAOCGTTA ACTTGTCAAA CTTCCAAACG	
TGACCACTGA ATGTAGCCAG GACCTTTTGT TCTTTCTGTG TGCCATGTAT GCCCCATTT	600
ACTGGTGACT TACATCGGTC CTGGAAAACA AGAAAGACAC ACGGTACATA CGGGGGTAAA	
GTACCATCGA TTTCAGCAT GAACCAATTA AGCCTTGCAA GTCOGTGTGC GAAAGGGCCA	660
CATGGTAGCT AAAGGTGTA CTTGGTTAAT TCGGAAOGTT CAGGCACACG CTTTCCCGGT	
GGGCGGCTG TGAGCCCAT CTCATAAAGT ACCGGCACAC TTGGCCAGAG AGCCTGGCAT	720
CCGGCCGAC ACTCGGGTAA GAGTATTTCA TGGCCGTGTG AACCAGTCTC TCGGACCGTA	
GTGAAGAGCT GCCCGTATAT GACAGAGGAG TCTGCATCTC CCCAGAGGCT ATCGTCACAG	780
CACTTCTCGA CGGGCATATA CTGTCTCTC AGACGTAGAG GGGTCTCCGA TAGCAGTGTC	
TGGAACAAGG AACAGATTCA ATGCCAGACT TCTCCATGGA TTCAAACAAT GGAAATTGCG	840
ACCTGTGTTT TTGTCTAAGT TACGGTCTGA AGAGGTACCT AAGTTTGTTA CCTTTAACGC	
GAAGCGGCAG GGAGCACTGT AAATGCAAGC CCATGAAGGC AACCCAAAAG ACGTATCTCA	900
CTTCGCGTC CCTCGTGACA TTTACGTTTC GGTACTTCCG TTGGGTTTTT TGCATAGAGT	
AGAATAATTA CAATTATGTA ATCAGAGCAA AAGTGAAAGA GGTGAAAGTG AAATGCCACG	960
TCTTATTAAT GTTAATACAT TAGTCTOGTT TTCACTTTCT CCACTTTCAC TTTACGGTGC	
ACGCAACAGC AATTGTGGAA GTAAAGGAGA TTCTCAAGTC TTCCCTAGTG AACATTCTTA	1020
TGCGTTGTGC TTAACACCTT CATTTCCTCT AAGAGTTCAG AAGGGATCAC TTGTAAGGAT	

Figure 4A

SUBSTITUTE SHEET (RULE 26)

AAGACACAGT TTCTGTGTCA	GACACTGTAC CTGTGACATG	ACCAACTCAG TGGTTGAGTC	GCTGCTTGTG CGACGAACAC	CCCCCAGCTT GGGGGTCGAA	GTTGCCAATG CAACGGTTAC	1080
AGGAATACAT TCCTTATGTA	AATTATGGGC TTAATACCCG	TATGAAGACA ATACTTCTGT	AAGAGCGTAC TTCTCGCATG	CAGGCTTCTA GTCOGAAGAT	CTAGTGGAAG GATCACCTTC	1140
GATCCTTGGC CTAGGAACCG	CGAAAAATGG GCTTTTTTACC	AGAGATCGTC TCTCTAGCAG	TTGCTAAGAA AACGATTCTT	AGTCAAGCGC TCAGTTCGCG	TGGGATCAAA ACCCCTAGTTT	1200
AGCTTCGACG TCGAAGCTGC	TCCCAGGAAA AGGGTCCTTT	AGCAAAGACC TCGTTTCTGG	CCGTGGCTCC GGCACCAGAG	AATTCCCAAC TTAAGGGTTG	AAAAACAGCA TTTTTGTCGT	1260
ATTCCAGACA TAAGGTCTGT	AGCGCGTAGT TCGCGCATCA	TAGACTAACG ATCTGATTGC	GAAAGGTGTA CTTTCACAT	TGGAACTCT ACCTTTGAGA	ATGGACTTTG TACCTGAAAC	1320
AAACTAAGAT TTTGATTCTA	TTGCATTGTT AACGTAACAA	GGAAGAGCAA CCTTCTCGTT	AAAAGAAATT TTTTCTTTAA	GCACTACAGC CGTGATGTCG	ACGTTATATT TGCAATATAA	1380
CTATTGTTTA GATAACAAAT	CTACAAGAAG GATGTTCTTC	CTGGTTTAGT GACCAAATCA	TGATTGTAGT ACTAACATCA	TCTCCTTTCC AGAGGAAAGG	TTCTTTTTTT AAGAAAAAAA	1440
TTATAACTAT AATATTGATA	ATTTGCACGT TAAACGTGCA	GTTCCCAGGC CAAGGGTCCG	AATTGTTTTA TTAACAAAAT	TTCAACTTCC AAGTTGAAGG	AGTGACAGAG TCACTGTCTC	1500
CAGTGACTGA GTCACTGACT	ATGTCTCAGC TACAGAGTCG	CTAAGAAGC GATTTCTTCG	TCAATTCATT AGTTAAGTAA	TCTGATCAAC AGACTAGTTG	TAATGGTGAC ATTACCACTG	1560
AAGTGTGTTGA TTCACAAACT	TACTTGGGGA ATGAACCCCT	AAGTGAACTA TTCATTGAT	ATTGCAATGG TAACGTTACC	TAAATCAGAG ATTTAGTCTC	AAAAGTTGAC TTTTCAACTG	1620
CAATGTTGCT GTTACAACGA	TTTCCTGTAG AAAGGACATC	ATGAACAAGT TACTTGTTCA	GAGAGATCAC CTCTCTAGTG	ATTTAAATGA TAAATTTACT	TGATCACTTT ACTAGTGAAA	1680
CCATTTAATA GGTAAATTAT	CTTTCAGCAG GAAAGTCGTC	TTTTAGTTAG AAAATCAATC	ATGACATGTA TACTGTACAT	GGATGCACCT CCTACGTGGA	AAATCTAAAT TTTAGATTTA	1740
ATTTTATCAT TAAAATAGTA	AAATGAAGAG TTTACTTCTC	CTGGTTTAGA GACCAAATCT	CTGTATGGTC GACATAACCAG	ACTGTTGGGA TGACAAOCT	AGGTAAATGC TCCATTTACG	1800
CTACTTTGTC GATGAAACAG	AATTCTGTTT TTAAGACAAA	TAAAAATTGC ATTTTTAACG	CTAAATAAAT GATTTATTTA	ATTAAGTCCT TAATTCAGGA	AAATAAAAAA TTTATTTTTT	1860
AAAAA TTTTTTTTT						

Figure 4B
SUBSTITUTE SHEET (RULE 26)

MLLLFRAIPM LLLGLMVLQT DCEIAQYYID EEEPPGTVIA VLSQHSIFNT TDIPATNFRL	60
MKQFNNSLIG VRESQGQLSI MERIDREQIC RQSLHCNLAL DVVSFSKGFH KLLNVKVEVR	120
DINDHSPHFP SEIMHVEVSE SSSVGTRIPL EIAIDEDVGS NSIQNFQISN NSHFSIDVLT	180
RADGVKYADL VLMRELDREI QPTYIMELLA MDGGVPSLSG TAVVNIRVLD FNDNSPVFER	240
STIAVDLVED APLGYLLEL HATDDDEGVN GEIVYGFSTL ASQEVRLFK INSRTGSVTL	300
EGQVDFETKQ TYEFEVQAQD LGPNPLTATC KVTVHILDVN DNTPAITITP LTTVNAGVAY	360
IPETATKENF IALISTTDRA SGSNGQVRCT LYGHEHFKLQ QAYEDSYMIV TTSTLDRENI	420
AAYSLTVVAE DLGFPSLTKK KYITVKVSD E NDNAPVFSKP QYEASILENN APGSYITTVI	480
ARDSDSQNG KVNRYRLVDAK VMGQSLTTFV SLDADSGVLR AVRSIDYEKL KQLDFEIEAA	540
DNGIPQLSTR VQLNLRIVDQ NDNCPVITNP LLNNGSGEVL LPISAPQNYL VFQLKAEDSD	600
EGHNSQLFYT ILRDP SRLFA INKESGEVFL KKQLNSDHSE DLSIVVAVYD LGRPSLSTNA	660
TVKFILTSF PSNVEVILQ PSAEEQHQID MSIIFI AVL GGCALLLAI FTVACTCKKK	720
AGEFKQVPEQ HGTCNEERLL STPSPQSVSS SLSQSESCQL SINTESENC S VSSNQEQHQ	780
TGIKHSISVP SYHTSGWHL D NCAM SISGHS HMGHISTKVQ WAKEIVTSMT VTLILVENQK	840
RRALSSQCRH KPVLTQMNQ QGSDMPITIS ATESTRVQKM GTAHCNMKRA IDCLTL	

Figure 5
SUBSTITUTE SHEET (RULE 26)

GAATCCCAG	AGATGAACTC	CTTGAGATTG	TTTTAATGA	CTGCAGGTCT	GGAAGGATTC	60
CTTAAGGGTC	TCTACTTGAG	GAAGCTCTAAC	AAAATTTACT	GACGTCCAGA	CCTTCCTAAG	
ACATTGCCAC	ACTGTTTCTA	GGCATGAAAA	AACTGCAAGT	TTCAACTTTG	TTTTTGGTGC	120
TGTAACGGTG	TGACAAAGAT	CCGTACTTTT	TTGACGTTCA	AAGTTGAAAC	AAAAACCACG	
AACTTTGATT	CTTCAAGATG	CTGCTTCTCT	TCAGAGCCAT	TCCAATGCTG	CTGTTGGGAC	180
TTGAAACTAA	GAAGTTCTAC	GACGAAGAGA	AGTCTCGGTA	AGGTTACGAC	GACAACCCTG	
TGATGGTTTT	ACAAACAGAC	TGTGAAATTG	CCCAGTACTA	CATAGATGAA	GAAGAACCCC	240
ACTACCAAAA	TGTTTGTCTG	ACACTTTAAC	GGGTACATGAT	GTATCTACTT	CTTCTTGGGG	
CTGGCACTGT	AATTGCAGTG	TTGTCACAAC	ACTCCATATT	TAACACTACA	GATATACCTG	300
GACCGTGACA	TTAACGTCAC	AACAGTGTTG	TGAGGTATAA	ATTGTGATGT	CTATATGGAC	
CAACCAATTT	CCGTCTAATG	AAGCAATTTA	ATAATTCCCT	TATCGGAGTC	CGTGAGAGTG	360
GTTGGTTAAA	GGCAGATTAC	TTCGTTAAAT	TATTAAGGGA	ATAGCCTCAG	GCACTCTCAC	
ATGGGCAGCT	GAGCATCATG	GAGAGGATTG	ACCGGGAGCA	AATCTGCAGG	CAGTCCCTTC	420
TACCCGTCGA	CTCGTAGTAC	CTCTCCTAAC	TGGCCCTCGT	TTAGACGTCC	GTCAGGGAAG	
ACTGCAACCT	GGCTTTGGAT	GTGGTCAGCT	TTTCCAAAGG	ACACTTCAAG	CTTCTGAACG	480
TGACGTTGGA	CCGAAACCTA	CACCAGTCGA	AAAGGTTTCC	TGTGAAGTTC	GAAGACTTGC	
TGAAAGTGGA	GGTGAGAGAC	ATTAATGACC	ATAGCCCTCA	CTTTCCAGT	GAAATAATGC	540
ACTTTCACCT	CCACTCTCTG	TAATTACTGG	TATCGGGAGT	GAAAGGGTCA	CTTTATTACG	
ATGTGGAGGT	GTCTGAAAGT	TCCTCTGTGG	GCACCAGGAT	TCCTTTAGAA	ATTGCAATAG	600
TACACCTCCA	CAGACTTTCA	AGGAGACACC	CGTGGTCCTA	AGGAAATCTT	TAACGTTATC	
ATGAAGATGT	TGGGTCCAAC	TOCATCCAGA	ACTTTCAGAT	CTCAAATAAT	AGCCACTTCA	660
TACTTCTACA	ACCCAGGTTG	AGGTAGGTCT	TGAAAGTCTA	GAGTTTATTA	TCGGTGAAGT	
GCATTGATGT	GCTAACCAGA	GCAGATGGGG	TGAAATATGC	AGATTTAGTC	TTAATGAGAG	720
CGTAACTACA	CGATTGGTCT	CGTCTACCCC	ACTTTATACG	TCTAAATCAG	AATTACTCTC	
AACTGGACAG	GGAAATCCAG	CCAACATACA	TAATGGAGCT	ACTAGCAATG	GATGGGGGTG	780
TTGACCTGTC	CCTTTAGGTC	GGTTGTATGT	ATTACCTCGA	TGATCGTTAC	CTACCCOCAC	
TACCATCACT	ATCTGGTACT	GCAGTGGTTA	ACATCCGAGT	CCTGGACTTT	AATGATAACA	840
ATGGTAGTGA	TAGACCATGA	CGTCACCAAT	TGTAGGCTCA	GGACCTGAAA	TTACTATTGT	
GCCCAGTGTT	TGAGAGAAGC	ACCATTGCTG	TGGACCTAGT	AGAGGATGCT	CCTCTGGGAT	900
CGGGTCACAA	ACTCTCTTCG	TGGTAACGAC	ACCTGGATCA	TCTCCTACGA	GGAGACCCTA	
ACCTTTTGTT	GGAGTTACAT	GCTACTGACG	ATGATGAAGG	AGTGAATGGA	GAAATTGTTT	960
TGGAAAACAA	CCTCAATGTA	CGATGACTGC	TACTACTTCC	TCACTTACCT	CTTTAACAAA	
ATGGATTGAG	CACTTTGGCA	TCTCAAGAGG	TACGTCAGCT	ATTTAAAATT	AACTCCAGAA	1020
TACCTAAGTC	GTGAAACCGT	AGAGTTCTCC	ATGCAGTCGA	TAAATTTTAA	TTGAGGTCTT	

Figure 6A
SUBSTITUTE SHEET (RULE 26)

CTGGCAGTGT	TACTCTTGAA	GGCCAAGTTG	ATTTTGAGAC	CAAGCAGACT	TACGAATTTG	1080
GACCGTCACA	ATGAGAACTT	CCGGTTCAAC	TAAAACCTCTG	GTTCTGCTGA	ATGCTTAAAC	
AGGTACAAGC	CCAAGATTTG	GGCCCCAACC	CACTGACTGC	TACTTGTAAG	GTAAGTGTTC	1140
TCCATGTTTC	GGTTCTAAAC	CCGGGGTTGG	GTGACTGACG	ATGAACATTT	CATTGACAAG	
ATATACTTGA	TGTAAATGAT	AATACCCAG	CCATCACTAT	TACCCCTCTG	ACTACTGTAA	1200
TATATGAACT	ACATTTACTA	TTATGGGGTC	GGTAGTGATA	ATGGGGAGAC	TGATGACATT	
ATGCAGGAGT	TGCCTATATT	CCAGAAACAG	CCACAAAGGA	GAACCTTTATA	GCTCTGATCA	1260
TACGTCCTCA	ACGGATATAA	GGTCTTTGTC	GGTGTTCCT	CTTGAAATAT	CGAGACTAGT	
GCACTACTGA	CAGAGCCTCT	GGATCTAATG	GACAAGTTCG	CTGTACTCTT	TATGGACATG	1320
CGTGATGACT	GTCTCGGAGA	CCTAGATTAC	CTGTTCAAGC	GACATGAGAA	ATACCTGTAC	
AGCACTTTAA	ACTACAGCAA	GCTTATGAGG	ACAGTTACAT	GATAGTTACC	ACCTCTACTT	1380
TCGTGAAATT	TGATGTCGTT	CGAATACTCC	TGTCAATGTA	CTATCAATGG	TGGAGATGAA	
TAGACAGGGA	AAACATAGCA	GCGTACTCTT	TGACAGTAGT	TGCAGAAGAC	CTTGGCTTCC	1440
ATCTGTCCCT	TTTGTATCGT	CGCATGAGAA	ACTGTCATCA	ACGTCTTCTG	GAAACGAAGG	
CCTCATTGAA	GACCAAAAAG	TACTACACAG	TCAAGGTTAG	TGATGAGAAT	GACAATGCAC	1500
GGAGTAACTT	CTGGTTTTTC	ATGATGTGTC	AGTTCCAATC	ACTACTCTTA	CTGTTACGTG	
CTGTATTTTC	TAAACCCAG	TATGAAGCTT	CTATTCTGGA	AAATAATGCT	CCAGGCTCTT	1560
GACATAAAAAG	ATTTGGGGTC	ATACTTCGAA	GATAAGACCT	TTTATTACGA	GGTCCGAGAA	
ATATAACTAC	AGTGATAGCC	AGAGACTCTG	ATAGTGATCA	AAATGGCAA	GTAAATTACA	1620
TATATTGATG	TCACTATCGG	TCTCTGAGAC	TATCACTAGT	TTTACOGTTT	CATTTAATGT	
GACTTGTTGA	TGCAAAAGTG	ATGGGCCAGT	CACTAACAAC	ATTTGTTTCT	CTTGATGCGG	1680
CTGAACACCT	ACGTTTTTAC	TACCOGGTCA	GTGATTGTTG	TAAACAAAGA	GAAGTACGCC	
ACTCTGGAGT	ATTGAGAGCT	GTTAGGTCTT	TAGACTATGA	AAAACCTTAA	CAACTGGATT	1740
TGAGACCTCA	TAACTCTCGA	CAATCCAGAA	ATCTGATACT	TTTTGAATTT	GTTGACCTAA	
TTGAAATTGA	AGCTGCAGAC	AATGGGATCC	CTCAACTCTC	CACTCGCGTT	CAACTAAATC	1800
AACCTTAACT	TCGACGTCTG	TTACCOCTAGG	GAGTTGAGAG	GTGAGCGCAA	GTTGATTTAG	
TCAGAATAGT	TGATCAAAAT	GATAATTGOC	CTGTGATAAC	TAATCCTCTT	CCTAATAATG	1860
AGTCTTATCA	ACTAGTTTTA	CTATTAAACGG	GACACTATTG	ATTAGGAGAA	GAATTATTAC	
GCTCGGGTGA	AGTTCTGCTT	CCCATCAGCG	CTOCTCAAAA	CTATTTAGTT	TTCCAGCTCA	1920
CGAGCCCACT	TCAAGACGAA	GGGTAGTCGC	GAGGAGTTTT	GATAAATCAA	AAGGTCGAGT	
AAGCCGAGGA	TTCAGATGAA	GGGCACAAC	CCAGCTGTT	CTATACCATA	CTGAGAGATC	1980
TTGGGCTCCT	AAGTCTACTT	CCCGTGTGTA	GGGTCGACAA	GATATGGTAT	GACTCTCTAG	
CAAGCAGATT	GTTTGCCATT	AACAAAGAAA	GTGGTGAAGT	GTTCTGAAA	AAACAATTAA	2040
GTTOGTCTAA	CAAACGGTAA	TTGTTTCTTT	CACCACTTCA	CAAGGACTTT	TTTGTTAATT	
ACTCTGACCA	TTCAGAGGAC	TTGAGCATAG	TAGTTGCAGT	GTATGACTTG	GGAAGACCTT	2100
TGAGACTGGT	AAGTCTCCTG	AACCTCGTATC	ATCAACGTCA	CATACTGAAC	CCTTCTGGAA	
CATTATCCAC	CAATGCTACA	GTTAAATTCA	TCCTCACCAG	CTCTTTTCTT	TCTAACGTTG	2160
GTAATAGGTG	GTTACGATGT	CAATTTAAGT	AGGAGTGGCT	GAGAAAAGGA	AGATTGCAAC	

Figure 6B

SUBSTITUTE SHEET (RULE 26)

AAGTCGTTAT	TTTGCAACCA	TCTGCAGAAG	AGCAGCACCA	GATCGATATG	TCCATTATAT	2220
TTTCAGCAATA	AAACGTTGGT	AGACGTCTTC	TCGTCGTGGT	CTAGCTATAC	AGGTAATATA	
TCATTGCAGT	GCTGGCTGGT	GGTTGTGCTT	TGCTACTTTT	GGCCATCTTT	TTTGTGGCCT	2280
AGTAACGTCA	CGACCGACCA	CCAACACGAA	ACGATGAAAA	CCGGTAGAAA	AAACACCGGA	
GTACTTGTA	AAAGAAAGCT	GGTGAATTTA	AGCAGGTACC	TGAACAACAC	GGAACATGCA	2340
CATGAACATT	TTTCTTTCGA	CCACTTAAAT	TCGTCCATGG	ACTTGTTGTG	CCTTGTACGT	
ATGAAGAACG	CCTGTTAAGC	ACCCCATCTC	CCCAGTOGGT	CTCTTCTTCT	TTGTCTCAGT	2400
TACTTCTTGC	GGACAATTCT	TGGGGTAGAG	GGGTCAGCCA	GAGAAGAAGA	AACAGAGTCA	
CTGAGTCATG	CCAACTCTCC	ATCAATACTG	AATCTGAGAA	TTGCAGCGTG	TCCTCTAACC	2460
GACTCAGTAC	GGTTGAGAGG	TAGTTATGAC	TTAGACTCTT	AACGTCGCAC	AGGAGATTGG	
AAGAGCAGCA	TCAGCAAACA	GGCATAAAGC	ACTCCATCTC	TGTACCATCT	TATCACACAT	2520
TTCTCGTCGT	AGTCGTTTGT	CCGTATTTCT	TGAGGTAGAG	ACATGGTAGA	ATAGTGTGTA	
CTGGTTGGCA	CCTGGACAAT	TGTGCAATGA	GCATAAGTGG	ACATTCTCAC	ATGGGGCACA	2580
GACCAACCGT	GGACCTGTTA	ACACGTTACT	CGTATTCCAC	TGTAAGAGTG	TACCCCGTGT	
TTAGTACAAA	GGTACAGTGG	GCAAAGGAGA	TAGTGACTTC	AATGACAGTG	ACTCTGATAC	2640
AATCATGTTT	CCATGTCACC	CGTTTCCTCT	ATCACTGAAG	TTACTGTCAC	TGAGACTATG	
TAGTGGAGAA	TCAGAAAAGA	AGAGCATTGA	GCAGCCAATG	CAGGCACAAG	CCAGTGCTCA	2700
ATCACCTCTT	AGTCTTTTCT	TCTCGTAACT	CGTCGGTTAC	GTCCGTGTTC	GGTCACGAGT	
ATACACAGAT	GAATCAGCAG	GGTTCCGACA	TGCCGATAAC	TATTTTCAGCC	ACCGAATCAA	2760
TATGTGTCTA	CTTAGTCGTC	CCAAGGCTGT	ACGGCTATTG	ATAAAGTCGG	TGGCTTAGTT	
CAAGGGTCCA	GAAAATGGGA	ACTGCACATT	GCAATATGAA	AAGGGCTATA	GACTGTCTTA	2820
GTTCCCAGGT	CTTTTACCCT	TGACGTGTAA	CGTTATACTT	TTCCCGATAT	CTGACAGAAT	
CTCTGTAGCT	CCTGTATATT	ACAATACCTA	CCATGCAAGA	ATGCCTAACC	TGCACATACC	2880
GAGACATCGA	GGACATATAA	TGTTATGGAT	GGTACGTTCT	TACGGATTGG	ACGTGTATGG	
GAACCATAOC	CTTAGAGACC	CTTATTACCA	TATCAATAAT	CCTGTTGCTA	ATCGGATGCA	2940
CTTGGTATGG	GAATCTCTGG	GAATAATGGT	ATAGTTATTA	GGACAACGAT	TAGCCTACGT	
GGCGGAATAT	GAAAGAGATT	TAGTCAACAG	AAGTGCAACG	TTATCTCCGC	AGAGATCGTC	3000
CCGCCTTATA	CTTTCTCTAA	ATCAGTTGTC	TTCACGTTGC	AATAGAGGCG	TCTCTAGCAG	
TAGCAGATAC	CAAGAATTCA	ATTACAGTCC	GCAGATATCA	AGACAGCTTC	ATCCTTCAGA	3060
ATCGTCTATG	GTTCTTAAGT	TAATGTCAGG	CGTCTATAGT	TCTGTGGAAG	TAGGAAGTCT	
AATTGCTACA	ACCTTTTAAT	CATTAGGCAT	GCAAGTGAGA	ATGCACAAAG	GCAAGTGCTT	3120
TTAACGATGT	TGGAAAATTA	GTAATCOGTA	CGTTCACTCT	TACGTGTTTC	CGTTCAAGAA	
TAGCATGAAA	GCTAAATATA	TGGAGTCTCC	CCTTTCCCTC	TGATGGATGG	GGGGAGACAC	3180
ATCGTACTTT	CGATTTATAT	ACCTCAGAGG	GGAAAGGGAG	ACTACCTACC	CCCCCTGTGT	
AGGACAGTGC	ATAAATATAC	AGCTGCTTTC	TATTTGCATT	TCACTTGGGA	ATTTTTTGTT	3240
TCCTGTCAAG	TATTTATATG	TCGACGAAAG	ATAAACGTAA	AGTGAACCCCT	TAAAAACAA	
TTTTTTACAT	ATTTATTTTT	CCTGAATTGA	ATGTGACATT	GTCCTGTCAC	CTAACTAGCA	3300
AAAAATGTA	TAAATAAAAA	GGACTTAACT	TACACTGTAA	CAGGACAGTG	GATTGATCGT	

Figure 6C
SUBSTITUTE SHEET (RULE 26)

ATTAAATCCA	CAGACCTACA	GTCAAATATT	TGAGGGCCCC	TGAAACAGCA	CATCAGTCAG	3360
TAATTTAGGT	GTCTGGATGT	CAGTTTATAA	ACTCCCCGGG	ACTTTGTCGT	GTAGTCAGTC	
GACCTAAAGT	GGCCTTTTTA	CTTTTAGCAG	CTCCTGGGTC	TGCCCTCTGT	GTTAATCAGC	3420
CTGGATTTCA	CCGGAAAAAT	GAAAATCGTC	GAGGACCCAG	ACGGGAGACA	CAATTAGTCG	
CCCTGGTCAA	GTCCTGAGTA	GGATCATGGC	GTTTTTATAT	GCATCTCACC	TACTTTGGAC	3480
GGGACCAGTT	CAGGACTCAT	CCTAGTACCG	CAAAAATATA	CGTAGAGTGG	ATGAAACCTG	
GTGATTTACA	CATAATAGGA	AACGCTTGGT	TTCAGTGAAG	TCTGTGTTGT	ATATATTCTG	3540
CACTAAATGT	GTATTATCCT	TTGCGAACCA	AAGTCACTTC	AGACACAACA	TATATAAGAC	
TTATATACAC	GCATTTTGTG	TTTGTGTATA	TATTTCAAGT	CCATTCAGAT	ATGTGTATAT	3600
AATATATGTG	CGTAAACAC	AAACACATAT	ATAAAGTTCA	GGTAAGTCTA	TACACATATA	
AGTGCAGACC	TTGTAAATTA	AATATTCTGA	TACTTTTTCC	TCAATAAATA	TTTAAAT	
TCACGTCTGG	AACATTTAAT	TTATAAGACT	ATGAAAAAGG	AGTTATTTAT	AAATTTA	

Figure 6D
SUBSTITUTE SHEET (RULE 26)

MVCCGPGRML LGWAGLLVLA ALCLLQVPGA QAAACEPVRI PLCKSLPWNM TKMPNHLHHS	60
TQANAILAME QFEGLLGTHC SPDLLFFLCA MYAPICTIDF QHEPIKPCKS VCERARQGCE	120
PILIKYRHSW PESLACDELP VYDRGVCISP EAIVTADGAD FPMDSSTGHC RGASSERCKC	180
KPV RATQKTY FRNNYNYVIR AKVKEVKMKC HDVTAVVEVK EILKASLVNI PRDTVNL YTT	240
SGCLCPPLTV NEEYVIMGYE DEERSRLLLV EGSIAEKWKD RLGKKVKRWD MKLRHLGLGK	300
TDASDSTQNNQ KSGRNSNPRP ARS.	

AAGCCTGGGA	CCATGGTCTG	CTGCGGCCCCG	GGACGGATGC	TGCTAGGATG	GGCCGGGTTG	60
TTCGGACCCT	GGTACCAGAC	GACGCCGGGC	CCTGCCTACG	ACGATCCTAC	CCGGCCCAAC	
CTAGTCCTGG	CTGCTCTCTG	CCTGCTCCAG	GTGCCCCGAG	CTCAGGCTGC	AGCCTGTGAG	120
GATCAGGACC	GACGAGAGAC	GGACGAGGTC	CACGGGCCCTC	GAGTCCGACG	TCCGACACTC	
CCTGTCCGCA	TCCCGCTGTG	CAAGTCCCTT	CCCTGGAACA	TGACCAAGAT	GCCCAACCAC	180
GGACAGGCGT	AGGGCGACAC	GTTCAGGGAA	GGGACCTTGT	ACTGGTTCTA	CGGGTTGGTG	
CTGCACCACA	GCACCCAGGC	TAACGCCATC	CTGGCCATGG	AACAGTTCGA	AGGGCTGCTG	240
GACGTGGTGT	CGTGGGTCCG	ATTGCGGTAG	GACCGGTACC	TTGTCAAGCT	TCCCGACGAC	
GGCACCCACT	GCAGCCCGGA	TCTTCTCTTC	TTCTCTGTG	CAATGTACGC	ACCCATTTGC	300
CCGTGGGTGA	CGTCGGGCCT	AGAAGAGAAG	AAGGAGACAC	GTTACATGCG	TGGGTAAACG	
ACCATCGACT	TCCAGCACGA	GCCCATCAAG	CCCTGCAAGT	CTGTGTGTGA	GCGCGCCCGA	360
TGGTAGCTGA	AGGTCGTGCT	CGGGTAGTTC	GGGACGTTCA	GACACACACT	C GCGCGGGCT	
CAGGGCTGCG	AGCCCATTTCT	CATCAAGTAC	CGCCACTCGT	GGCCGGAAAG	CTTGGCCTGC	420
GTCCCGACGC	TCGGGTAAGA	GTAGTTTCATG	GCGGTGAGCA	CCGGCCTTTC	GAACCGGACG	
GACGAGCTGC	CGGTGTACGA	CCGCGGCGTG	TGCATCTCTC	CTGAGGCCAT	CGTCACCGCG	480
CTGCTCGACG	GCCACATGCT	GGCGCCGCAC	ACGTAGAGAG	GACTCCGGTA	GCAGTGGCGC	
GACGGAGCGG	ATTTTCCTAT	GGATTCAAGT	ACTGGACACT	GCAGAGGGGC	AAGCAGCGAA	540
CTGCCTCGCC	TAAAAGGATA	CCTAAGTTCA	TGACCTGTGA	CGTCTCCCCG	TTCGTGCTTT	
CGTTGCAAAT	GTAAGCCTGT	CAGAGCTACA	CAGAAGACCT	ATTTCCGGAA	CAATTACAAC	600
GCAACGTTTA	CATTCCGACA	GTCTCGATGT	GTCTTCTGGA	TAAAGGCCTT	GTTAATGTTG	
TATGTCATCC	GGGCTAAAGT	TAAAGAGGTA	AAGATGAAAT	GTCATGATGT	GACCGCCGTT	660
ATACAGTAGG	CCCGATTTC	ATTTCTCCAT	TTCTACTTTA	CAGTACTACA	CTGGCGGCAA	
GTGGAAGTGA	AGGAAATTCT	AAAGGCATCA	CTGGTAAACA	TTCCAAGGGA	CACCGTCAAT	720
CACCTTCACT	TCCTTTAAGA	TTTCCGTAGT	GACCATTGTG	AAGGTTCCCT	GTGGCAGTTA	
CTTTATACCA	CCTCTGGCTG	CCTCTGTCCT	CCACTTACTG	TCAATGAGGA	ATATGTCATC	780
GAAATATGGT	GGAGACCGAC	GGAGACAGGA	GGTGAATGAC	AGTTACTCCT	TATACAGTAG	
ATGGGCTATG	AAGACGAGGA	ACGTTCCAGG	TTACTCTTGG	TAGAAGGCTC	TATAGCTGAG	840
TACCCGATAC	TTCTGCTCCT	TGCAAGGTCC	AATGAGAACC	ATCTTCCGAG	ATATCGACTC	
AAGTGGAAGG	ATCGGCTTGG	TAAGAAAGTC	AAGCGCTGGG	ATATGAAACT	CCGACACCTT	900
TTCACCTTCC	TAGCCGAACC	ATTCTTTTCAG	TTGCGGACCC	TATACTTTGA	GGCTGTGGAA	
GGACTGGGTA	AAACTGATGC	TAGCGATTCC	ACTCAGAATC	AGAAGTCTGG	CAGGAACTCT	960
CCTGACCCAT	TTTGACTACG	ATCGCTAAGG	TGAGTCTTAG	TCTTCAGACC	GTCTTGAGA	

Figure 8A
SUBSTITUTE SHEET (RULE 26)

AATCCCCGGC	CAGCACGCAG	CTAAATCCTG	AAATGTAAAA	GGCCACACCC	ACGGACTCCC	1020
TTAGGGGCCG	GTCGTGCGTC	GATTTAGGAC	TTTACATTTT	CCGGTGTGGG	TGCCTGAGGG	
TTCTAAGACT	GGCGCTGGTG	GACTAACAAA	GGAAAACCGC	ACAGTTGTGC	TCGTGACCGA	1080
AAGATTCTGA	CCGCGACCAC	CTGATTGTTT	CCTTTTGGCG	TGTCAACACG	AGCACTGGCT	
TTGTTTACCG	CAGACACCGC	GTGGCTACCG	AAGTTACTTC	CGGTCCCCTT	TCTCCTGCTT	1140
AACAAATGGC	GTCTGTGGCG	CACCGATGGC	TTCAATGAAG	GCCAGGGGAA	AGAGGACGAA	
CTTAATGGCG	TGGGGTTAGA	TCCTTTAATA	TGTTATATAT	TCTGTTTCAT	CAATCACGTG	1200
GAATTACCGC	ACCCCAATCT	AGGAAATTAT	ACAATATATA	AGACAAAGTA	GTTAGTGCAC	
GGGACTGTTC	TTTGTCAACC	AGAATAGTAA	ATTAAATATG	TTGATGCTAA	GGTTTCTGTA	1260
CCCTGACAAG	AAAACGTTGG	TCTTATCATT	TAATTTATAC	AAC TACGATT	CCAAAGACAT	
CTGGACTCCC	TGGGTTTAAT	TTGGTGTTC	GTACCCTGAT	TGAGAATGCA	ATGTTTCATG	1320
GACCTGAGGG	ACCCAAATTA	AACCACAAGA	CATGGGACTA	ACTCTTACGT	TACAAAGTAC	
TAAAGAGAGA	ATCCTGGTCA	TATCTCAAGA	ACTAGATATT	GCTGTAAGAC	AGCCTCTGCT	1380
ATTTCTCTCT	TAGGACCAGT	ATAGAGTTCT	TGATCTATAA	CGACATTCTG	TCGGAGACGA	
GCTGCGCTTA	TAGTCTTG TG	TTTGTATGCC	TTTGTCCATT	TCCCTCATGC	TGTGAAAGTT	1440
CGACGCGAAT	ATCAGAACAC	AAACATACGG	AAACAGGTAA	AGGGAGTACG	ACACTTTCAA	
ATACATGTTT	ATAAAGGTAG	AACGGCATT	TGAAATCAGA	CACTGCACAA	GCAGAGTAGC	1500
TATGTACAAA	TATTTCCATC	TTGCCGTAAA	ACTTTAGTCT	GTGACGTGTT	CGTCTCATCG	
CCAACACCAG	GAAGCATTTA	TGAGGAAACG	CCACACAGCA	TGACTTATTT	TCAAGATTGG	1560
GGTTGTGGTC	CTTCGTAAAT	ACTCCTTTGC	GGTGTGTCGT	ACTGAATAAA	AGTTCTAACC	
CAGGCAGCAA	AATAAATAGT	GTTGGGAGCC	AAGAAAAGAA	TATTTTGCCT	GGTTAAGGGG	1620
GTCCGTCGTT	TTATTTATCA	CAACCCTCGG	TTCTTTTCTT	ATAAAACGGA	CCAATTCCCC	
CACACTGGAA	TCAGTAGCCC	TTGAGCCATT	AACAGCAGTG	TTCTTCTGGC	AAGTTTTTGA	1680
GTGTGACCTT	AGTCATCGGG	AACTCGGTAA	TTGTCGTCAC	AAGAAGACCG	TTCAAAAAC	
TTTGTTTATA	AATGTATTCA	CGAGCATTAG	AGATGAACTT	ATAACTAGAC	ATCTGTTGTT	1740
AAACAAGTAT	TTACATAAGT	GCTCGTAATC	TCTACTTGAA	TATTGATCTG	TAGACAACAA	
ATCTCTATAG	CTCTGCTTCC	TTCTAAATCA	AACCCATTGT	TGGATGCTCC	CTCTCCATTC	1800
TAGAGATATC	GAGACGAAGG	AAGATTTAGT	TTGGGTAAAC	ACCTACGAGG	GAGAGGTAAG	

Figure 8B
SUBSTITUTE SHEET (RULE 26)

ATAAATAAAT	TTGGCTTGCT	GTATTGGCCA	GGAAAAGAAA	GTATTAAAGT	ATGCATGCAT	1860
TATTTATTTA	AACCGAACGA	CATAACCGGT	CCTTTTCTTT	CATAATTTCA	TACGTACGTA	
GTGCACCAGG	GTGTTATTTA	ACAGAGGTAT	GTAACCTCTAT	AAAAGACTAT	AATTTACAGG	1920
CACGTGGTCC	CACAATAAAT	TGTCTCCATA	CATTGAGATA	TTTCTTGATA	TTAAATGTCC	
ACACGGAAAT	GTGCACATTT	GTTTACTTTT	TTTCTTCCTT	TTGCTTTGGG	CTTGTGATTT	1980
TGTGCCTTTA	CACGTGTAAA	CAAATGAAAA	AAAGAAGGAA	AACGAAACCC	GAACACTAAA	
TGGTTTTTGG	TGTGTTTATG	TCTGTATTTT	GGGGGGTGGG	TAGGTTTAAG	CCATTGCACA	2040
ACCAAAAACC	ACACAAATAC	AGACATAAAA	CCCCCACC	ATCCAAATTC	GGTAACGTGT	
TTCAAGTTGA	ACTAGATTAG	AGTAGACTAG	GCTCATTGGC	CTAGACATTA	TGATTTGAAT	2100
AAGTTCAACT	TGATCTAATC	TCATCTGATC	CGAGTAACCG	GATCTGTAAT	ACTAAACTTA	
TTGTGTTGTT	TAATGCTCCA	TCAAGATGTC	TAATAAAAGG	AATATGGTTG	TCAACAGAGA	2160
AACACAACAA	ATTACGAGGT	AGTTCTACAG	ATTATTTTCC	TTATACCAAC	AGTTGTCTCT	
CGACAACAAC	AACAAA					
GCTGTTGTTG	TTGTTT					

Figure 8C
SUBSTITUTE SHEET (RULE 26)

MVCGSPGGML LLRAGLLALA ALCLLRVPGA RAAACEPVRI PLCKSLPWNM TKMPNHLHHS	60
TQANAILAIE QFEGLLGTHC SPDLLFFLCA MYAPICTIDF QHEPIKPCKS VCERARQGCE	120
PILIKYRHSW PENLACEELP VYDRGVCISP EAIVTADGAD FPMDSNGNC RGASSERCKC	180
KPIRATQKTY FRNNYNYVIR AKVKEIKTKC HDVTAVVEVK EILKSSLVNI PRDTVNLYTS	240
SGCLCPPLNV NEEYIIMGYE DEERSRLLL V EGSIAEKWKD RLGKKVKRWD MKLRHLGLSK	300
SDSSNSDSTQ SQKSGRNSNP RQARN.	

Figure 9
SUBSTITUTE SHEET (RULE 26)

GGCGGAGCGG	GCCTTTTGGC	GTCCACTGCG	CGGCTGCACC	CTGCCCCATC	TGCCGGGATC	60
CCGCCTCGCC	CGGAAAACCG	CAGGTGACGC	GCCGACGTGG	GACGGGGTAG	ACGGCCCTAG	
ATGGTCTGCG	GCAGCCCGGG	AGGGATGCTG	CTGCTGCGGG	CCGGGCTGCT	TGCCCTGGCT	120
TACCAGACGC	CGTCGGGCCC	TCCCTACGAC	GACGACGCC	GGCCCGACGA	ACGGGACCGA	
GCTCTCTGCC	TGCTCCGGGT	GCCCGGGGCT	CGGGCTGCAG	CCTGTGAGCC	CGTCCGCATC	180
CGAGAGACGG	ACGAGGCCCA	CGGGCCCCGA	GCCCGACGTC	GGACACTCGG	GCAGGCGTAG	
CCCCTGTGCA	AGTCCCTGCC	CTGGAACATG	ACTAAGATGC	CCAACCACCT	GCACCACAGC	240
GGGGACACGT	TCAGGGACGG	GACCTTGTAC	TGATTCTACG	GGTTGGTGGA	CGTGGTGTCTG	
ACTCAGGCCA	ACGCCATCCT	GGCCATCGAG	CAGTTCGAAG	GTCTGTGTTGG	CACCCACTGC	300
TGAGTCCGGT	TGCGGTAGGA	CCGGTAGCTC	GTCAAGCTTC	CAGACGACCC	GTGGGTGACG	
AGCCCCGATC	TGCTCTTCTT	CCTCTGTGCC	ATGTACGCGC	CCATCTGCAC	CATTGACTTC	360
TCGGGGCTAG	ACGAGAAGAA	GGAGACACGG	TACATGCGCG	GGTAGACGTG	GTAACCTGAAG	
CAGCACGAGC	CCATCAAGCC	CTGTAAGTCT	GTGTGCGAGC	GGGCCCCGCA	GGGCTGTGAG	420
GTCGTGCTCG	GGTAGTTCCG	GACATTGAGA	CACACGCTCG	CCCGGGCCGT	CCCGACACTC	
CCCATACTCA	TCAAGTACCG	CCACTCGTGG	CCGGAGAACC	TGGCCTGCGA	GGAGCTGCCA	480
GGGTATGAGT	AGTTCATGGC	GGTGAGCACC	GGCCTCTTGG	ACCGGACGCT	CCTCGACGGT	
GTGTACGACA	GGGGCGTGTG	CATCTCTCCC	GAGGCCATCG	TTACTGCGGA	CGGAGCTGAT	540
CACATGCTGT	CCCCGCACAC	GTAGAGAGGG	CTCCGGTAGC	AATGACGCCT	GCCTCGACTA	
TTTCCTATGG	ATTCTAGTAA	CGGAAACTGT	AGAGGGGCAA	GCAGTGAACG	CTGTAAATGT	600
AAAGGATACC	TAAGATCATT	GCCTTTGACA	TCTCCCCGTT	CGTCACTTGC	GACATTTACA	
AAGCCTATTA	GAGCTACACA	GAAGACCTAT	TTCCGGAACA	ATTACAACCTA	TGTCATTCCG	660
TTCGGATAAT	CTCGATGTGT	CTTCTGGATA	AAGGCCTTGT	TAATGTTGAT	ACAGTAAGCC	
GCTAAAGTTA	AAGAGATAAA	GACTAAGTGC	CATGATGTGA	CTGCAGTAGT	GGAGGTGAAG	720
CGATTTCAAT	TTCTCTATTT	CTGATTACAG	GTACTACACT	GACGTCATCA	CCTCCACTTC	
GAGATTCTAA	AGTCCTCTCT	GGTAAACATT	CCACGGGACA	CTGTCAACCT	CTATACCAGC	780
CTCTAAGATT	TCAGGAGAGA	CCATTTGTAA	GGTGCCCTGT	GACAGTTGGA	GATATGGTCG	
TCTGGCTGCC	TCTGCCCTCC	ACTTAATGTT	AATGAGGAAT	ATATCATCAT	GGGCTATGAA	840
AGACCGACGG	AGACGGGAGG	TGAATTACAA	TTACTCCTTA	TATAGTAGTA	CCCGATACTT	

Figure 10A
SUBSTITUTE SHEET (RULE 26)

GATGAGGAAC	GTTCCAGATT	ACTCTTGGTG	GAAGGCTCTA	TAGCTGAGAA	GTGGAAGGAT	900
CTACTCCTTG	CAAGGTCTAA	TGAGAACCAC	CTTCCGAGAT	ATCGACTCTT	CACCTTCCTA	
CGACTCGGTA	AAAAAGTTAA	GCGCTGGGAT	ATGAAGCTTC	GTCATCTTGG	ACTCAGTAAA	960
GCTGAGCCAT	TTTTTCAATT	CGCGACCCTA	TACTTCGAAG	CAGTAGAACC	TGAGTCATTT	
AGTGATTCTA	GCAATAGTGA	TTCCACTCAG	AGTCAGAAGT	CTGGCAGGAA	CTCGAACCCC	1020
TCACTAAGAT	CGTTATCACT	AAGGTGAGTC	TCAGTCTTCA	GACCGTCCTT	GAGCTTGGGG	
CGGCAAGCAC	GCAACTAAAT	CCCGAAATAC	AAAAAGTAAC	ACAGTGGACT	TCCTATTAAAG	1080
GCCGTTCGTG	CGTTGATTTA	GGGCTTTATG	TTTTTCATTG	TGTCACCTGA	AGGATAATTC	
ACTTACTTGC	ATTGCTGGAC	TAGCAAAGGA	AAATTGCACT	ATTGCACATC	ATATTCTATT	1140
TGAATGAACG	TAACGACCTG	ATCGTTTCCT	TTTAACGTGA	TAACGTGTAG	TATAAGATAA	
GTTTACTATA	AAAATCATGT	GATAACTGAT	TATTACTTCT	GTTTCTCTTT	TGGTTTCTGC	1200
CAAATGATAT	TTTTAGTACA	CTATTGACTA	ATAATGAAGA	CAAAGAGAAA	ACCAAAGACG	
TTCTCTCTTC	TCTCAACCCC	TTTGTAATGG	TTTGGGGGCA	GACTCTTAAG	TATATTGTGA	1260
AAGAGAGAAG	AGAGTTGGGG	AAACATTACC	AAACCCCCGT	CTGAGAAATC	ATATAACACT	
GTTTCTTATT	TCACTAATCA	TGAGAAAAAC	TGTTCTTTTG	CAATAATAAT	AAATTAAACA	1320
CAAAAGATAA	AGTGATTAGT	ACTCTTTTTG	ACAAGAAAAC	GTTATTATTA	TTTAATTTGT	
TGCTGTTACC	AGAGCCTCTT	TGCTGAGTCT	CCAGATGTTA	ATTTACTTTT	TGCACCCCAA	1380
ACGACAAATG	TCTCGGAGAA	ACGACTCAGA	GGTCTACAAT	TAAATGAAAG	ACGTGGGGTT	
TTGGGAATGC	AATATTGGAT	GAAAAGAGAG	GTTTCTGGTA	TTACACAGAA	GCTAGATATG	1440
AACCTTTACG	TTATAACCTA	CTTTTCTCTC	CAAAGACCAT	AAGTGTCTTT	CGATCTATAC	
CCTTAAACA	TACTCTGCCG	ATCTAATTAC	AGCCTTATTT	TTGTATGCCT	TTTGGGCATT	1500
GGAATTTTGT	ATGAGACGGC	TAGATTAATG	TCGGAATAAA	AACATACGGA	AAACCCGTAA	
CTCCTCATGC	TTAGAAAGTT	CCAAATGTTT	ATAAAGGTAA	AATGGCAGTT	TGAAGTCAAA	1560
GAGGAGTACG	AATCTTTCAA	GGTTTACAAA	TATTTCCATT	TTACCGTCAA	ACTTCAGTTT	
TGTCACATAG	GCAAAGCAAT	CAAGCACCAG	GAAGTGTTTA	TGAGGAAACA	ACACCCAAGA	1620
ACAGTGTATC	CGTTTCGTTA	GTTCGTGGTC	CTTCACAAAT	ACTCCTTTGT	TGTGGGTTCT	
TGAATTATTT	TTGAGACTGT	CAGGAAGTAA	AATAAATAGG	AGCTTAAGAA	AGAACATTTT	1680
ACTTAATAAA	AACTCTGACA	GTCCTTCATT	TTATTTATCC	TCGAATTCCT	TCTTGTAATA	
GCCTGATTGA	GAAGCACAA	TGAAACCAGT	AGCCGCTGGG	GTGTTAATGG	TAGCATTCCT	1740
CGGACTAACT	CTTCGTGTTG	ACTTTGGTCA	TCGGCGACCC	CACAATTACC	ATCGTAAGAA	
CTTTTGCCAA	TACATTTGAT	TTGTTTCATG	ATATATTAAT	CAGCATTAGA	GAAATGAATT	1800
GAAAACCGTT	ATGTAAACTA	AACAAGTACT	TATATAATTA	GTCGTAATCT	CTTTACTTAA	
ATAACTAGAC	ATCTGCTGTT	ATCACCATAG	TTTTGTTTAA	TTTGCTTCCT	TTTAAATAAA	1860
TATTGATCTG	TAGACGACAA	TAGTGGTATC	AAAACAAATT	AAACGAAGGA	AAATTTATTT	
CCCATTGGTG	AAAGTCAAAA	AAAAAAAAAA	AAA			
GGGTAACCAC	TTTCAGTTTT	TTTTTTTTTT	TTT			

Figure 10B
SUBSTITUTE SHEET (RULE 26)

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/10942**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) : Please See Extra Sheet.

US CL : 530/300, 350; 514/2; 536/23.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 530/300, 350; 514/2; 536/23.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

DIALOG (MEDLINE, BIOSIS, EMBASE, WPI, USPATFULL) AUTHOR AND WORD. search terms: e.g. cerberus, xenopus

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y, P	BOUWMEESTER et al. Cerberus is a head-inducing secreted factor expressed in the anterior endoderm of Spemann's organizer. Nature. 15 August 1996, Vol. 382, No. 6592, pages 595-601, see entire document.	1-15

☐ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

•	Special categories of cited documents:	T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
•A	document defining the general state of the art which is not considered to be of particular relevance		
•E	earlier document published on or after the international filing date	•X	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
•L	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	•Y	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
•O	document referring to an oral disclosure, use, exhibition or other means		
•P	document published prior to the international filing date but later than the priority date claimed	•A	document member of the same patent family

Date of the actual completion of the international search

29 AUGUST 1997

Date of mailing of the international search report

11 SEP 1997

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

HEATHER BAKALYAR

Telephone No. (703) 308-0196

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

A01N 37/18; A61K 38/00; C07K 1/00, 2/00, 4/00, 7/00, 14/00, 16/00, 17/00; C07H 21/02, 21/04

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.